

## Revelstoke IPMA Priority Plant List

<b>REGIONAL EDRR</b> - High risk invasive plant species not currently known within the Columbia Shuswap Regional District boundary or brand new incursions that are extremely limited in extent (less than 10 very small sites) within the Columbia Shuswap Regional District boundary. Management objective is eradication.		
<ul style="list-style-type: none"> <li>- Bighead knapweed</li> <li>- Buffalobur</li> <li>- Bur chervil</li> <li>- Colt's foot</li> <li>- Common bugloss</li> <li>- Field scabious</li> <li>- Garlic mustard</li> <li>- Giant hogweed</li> </ul>	<ul style="list-style-type: none"> <li>- Giant knotweed</li> <li>- Gorse</li> <li>- Greater knapweed</li> <li>- Himalayan knotweed</li> <li>- Hoary Cress</li> <li>- Japanese butterbur</li> <li>- Longspine sandbur</li> <li>- Nodding thistle</li> </ul>	<ul style="list-style-type: none"> <li>- North Africa grass</li> <li>- Plumeless thistle</li> <li>- Puncturevine</li> <li>- Short-fringed knapweed</li> <li>- Tansy Ragwort</li> <li>- Wild chervil</li> <li>- Wild parsnip</li> <li>- Wood sage</li> </ul>
<b>ERADICATION or ANNUAL CONTROL</b> – Species are known in the IPMA but with limited distribution and/or significant potential to spread. Management objective is to control sites annually.		
<ul style="list-style-type: none"> <li>- Baby's breath</li> <li>- Black knapweed (BC)</li> <li>- Blueweed</li> <li>- Bohemian knotweed</li> <li>- Himalayan blackberry</li> </ul>	<ul style="list-style-type: none"> <li>- Hoary Alyssum</li> <li>- Japanese knotweed</li> <li>- Leafy spurge (BC)</li> <li>- Marsh plume thistle</li> <li>- Poison hemlock</li> </ul>	<ul style="list-style-type: none"> <li>- Policeman's helmet</li> <li>- Scotch broom</li> <li>- Teasel</li> <li>- Yellow archangel</li> </ul>
<b>CONTAINMENT</b> – Species are abundant (with no expectation of eradication) in certain portions of the IPMA but have not yet infested all potential habitats. Containment is the management objective. Treat all sites outside of containment lines.		
<b>Contain to gardens:</b> <ul style="list-style-type: none"> <li>- Butterfly bush</li> <li>- Common periwinkle</li> <li>- English holly</li> <li>- English ivy</li> <li>- Garden yellow loosestrife</li> </ul>	<b>Contain to gardens Cont'd:</b> <ul style="list-style-type: none"> <li>- Goutweed</li> <li>- Mountain bluet</li> <li>- Russian olive</li> <li>- Salt cedar/ Tamarisk</li> <li>- Siberian elm</li> </ul>	<b>Contain to southern portion of IPMA:</b> <ul style="list-style-type: none"> <li>- Common tansy</li> </ul> <b>Contain to Revelstoke and Arrow Reservoirs:</b> <ul style="list-style-type: none"> <li>- Eurasian water milfoil</li> </ul>
<b>ESTABLISHED (BIOCONTROL OR SITE-SPECIFIC APPROACH)</b> – Widespread species beyond landscape-level control and/or have relatively low impact. May have biocontrol (BC) available. Treat based on land management objectives.		
<ul style="list-style-type: none"> <li>- Annual sow thistle</li> <li>- Bull thistle (BC)</li> <li>- Burdock</li> <li>- Canada thistle (BC)</li> <li>- Caraway</li> <li>- Chicory</li> <li>- Common comfrey</li> <li>- Curled dock</li> <li>- Dalmatian toadflax (BC)</li> </ul>	<ul style="list-style-type: none"> <li>- Diffuse knapweed (BC)</li> <li>- Hound's tongue (BC)</li> <li>- Knapweed spp. (BC)</li> <li>- Meadow buttercup</li> <li>- Meadow goat's beard</li> <li>- Meadow knapweed (BC)</li> <li>- Orange hawkweed</li> <li>- Oxeye daisy</li> <li>- Perennial sow thistle</li> </ul>	<ul style="list-style-type: none"> <li>- Purple loosestrife (BC)</li> <li>- Sow thistle spp.</li> <li>- Spotted knapweed (BC)</li> <li>- St. John's Wort (BC)</li> <li>- Sulphur cinquefoil</li> <li>- Western goat's beard</li> <li>- Wormwood</li> <li>- Yellow hawkweed spp.</li> <li>- Yellow toadflax (BC)</li> </ul>
<b>INSUFFICIENT INFORMATION</b> – Insufficient information for these species on their distribution, impacts, potential for spread and/or feasibility of control. Further information is required.		
<ul style="list-style-type: none"> <li>- Bachelor's button</li> <li>- Black locust</li> <li>- Brown knapweed</li> <li>- Carpet burweed</li> <li>- Creeping buttercup</li> <li>- Curly leaf pondweed</li> <li>- Cypress spurge</li> <li>- Dame's rocket</li> <li>- Eyebright</li> <li>- Field bindweed</li> </ul>	<ul style="list-style-type: none"> <li>- Flat peavine</li> <li>- Fragrant water lily</li> <li>- Greater celandine</li> <li>- Green foxtail</li> <li>- Hairy cat's ear</li> <li>- Hedge false bindweed</li> <li>- Kochia</li> <li>- Lady's thumb</li> <li>- Night-flowering catchfly</li> <li>- Nightshade</li> </ul>	<ul style="list-style-type: none"> <li>- Queen Anne's Lace</li> <li>- Rush skeletonweed (BC)</li> <li>- Russian knapweed</li> <li>- Russian thistle</li> <li>- Scentless chamomile (BC)</li> <li>- Scotch thistle</li> <li>- Spurge laurel</li> <li>- Sweet fennel</li> <li>- Wild four o'clock</li> <li>- Yellow flag iris</li> </ul>

BC – biocontrol