## Revelstoke IPMA Priority Plant List

PREVENT – High priority species not		ithin the region. Management objective is
	blishment. Bolded species indicate Provi	
<ul> <li>Bighead knapweed</li> <li>Black henbane</li> <li>Buffalobur</li> <li>Bur chervil</li> <li>Colt's foot</li> <li>Common bugloss</li> <li>Field scabious</li> <li>Giant hogweed</li> <li>Gorse</li> </ul>	<ul> <li>Himalayan knotweed</li> <li>Hoary cress</li> <li>Japanese butterbur</li> <li>Longspine sandbur</li> <li>Nodding thistle</li> <li>North Africa grass</li> <li>Plumeless thistle</li> <li>Poison hemlock</li> <li>Puncturevine</li> </ul>	<ul> <li>Rush skeletonweed (BC)</li> <li>Russian knapweed</li> <li>Scotch thistle</li> <li>Short-fringed knapweed</li> <li>Spurge laurel</li> <li>Tansy ragwort</li> <li>Tree of heaven</li> <li>Wood sage</li> <li>Yellow flag iris</li> </ul>
- Greater knapweed		
	ies extremely limited in extent (less thar . Management objective is eradication.	10 very small sites) within the Columbia
<ul><li>Cypress spurge</li><li>Garlic mustard</li></ul>	- Giant knotweed - Wild chervil hited extent and/ or significant potential	- Wild parsnip
	areas with the ultimate goal of reducing	
<ul> <li>Baby's breath</li> <li>Black knapweed (BC)</li> <li>Blueweed</li> <li>Bohemian knotweed</li> <li>Brown knapweed</li> </ul>	<ul> <li>Himalayan blackberry</li> <li>Hoary alyssum</li> <li>Japanese knotweed</li> <li>Leafy spurge (BC)</li> <li>Marsh plume thistle</li> </ul>	<ul> <li>Policeman's helmet</li> <li>Scotch broom</li> <li>Teasel</li> <li>Yellow archangel</li> </ul>
		anagement objective is to prevent further containment lines and identification of
occurrences outside the line to cont		containment lines and identification of
Contain to gardens: - Butterfly bush - Common periwinkle - English holly - English ivy - Garden yellow loosestrife	<ul> <li>Goutweed</li> <li>Mountain bluet</li> <li>Myrtle spurge</li> <li>Russian olive</li> <li>Salt cedar/ Tamarisk</li> <li>Siberian elm</li> </ul>	Contain to southern portion of IPMA: - Common tansy Contain to Revelstoke and Arrow Reservoirs: - Eurasian water milfoil
		ecific situations with certain high values -
		is to reduce the invasive species impacts
locally or regionally, where resource         -       Bull thistle (BC)         -       Burdock         -       Canada thistle (BC)         -       Caraway         -       Chicory         -       Common comfrey         -       Dalmatian toadflax (BC)         -       Diffuse knapweed (BC)	<ul> <li>Hound's tongue (BC)</li> <li>Meadow buttercup</li> <li>Meadow goat's beard</li> <li>Meadow knapweed (BC)</li> <li>Orange hawkweed</li> <li>Oxeye daisy</li> <li>Purple loosestrife (BC)</li> </ul>	<ul> <li>Spotted knapweed (BC)</li> <li>St. John's Wort (BC)</li> <li>Sulphur cinquefoil</li> <li>Western goat's beard</li> <li>Wormwood</li> <li>Yellow hawkweed spp.</li> <li>Yellow toadflax (BC)</li> </ul>
and/or feasibility of control. Further		distribution, impacts, potential for spread
<ul> <li>Bachelor's button</li> <li>Black locust</li> <li>Carpet burweed</li> <li>Cheatgrass/downy brome</li> <li>Creeping buttercup</li> <li>Curly leaf pondweed</li> <li>Dame's rocket</li> </ul>	<ul> <li>Field bindweed</li> <li>Flat peavine</li> <li>Fragrant water lily</li> <li>Greater celandine</li> <li>Green foxtail</li> <li>Kochia</li> <li>Lady's thumb</li> </ul>	<ul> <li>Nightshade</li> <li>Queen Anne's Lace</li> <li>Russian thistle</li> <li>Scentless chamomile (BC)</li> <li>Sweet fennel</li> <li>Wild four o'clock</li> <li>Woolly vetch</li> </ul>

BC – biocontrol