

## Salmon Arm IPMA Priority Plant List

<b>PREVENT</b> – High priority species not currently known in the IPMA and/ or within the region. Management objective is to prevent the introduction and establishment. <b>Bolded species indicate Provincial EDRR</b>		
<ul style="list-style-type: none"> <li>- Bighead knapweed</li> <li>- Black henbane</li> <li>- Buffalobur</li> <li>- Bur chervil</li> <li>- Garlic mustard</li> <li>- Giant knotweed</li> </ul>	<ul style="list-style-type: none"> <li>- Gorse</li> <li>- Greater knapweed</li> <li>- Himalayan knotweed</li> <li>- Longspine sandbur</li> <li>- Nodding thistle</li> <li>- North Africa grass</li> </ul>	<ul style="list-style-type: none"> <li>- Plumeless thistle</li> <li>- Puncturevine</li> <li>- Spurge laurel</li> <li>- Tansy ragwort</li> <li>- Wood sage</li> </ul>
<b>REGIONAL EDRR</b> – High priority species extremely limited in extent (less than 10 very small sites) within the Columbia Shuswap Regional District boundary. Management objective is eradication.		
<ul style="list-style-type: none"> <li>- Common bugloss</li> <li>- Cypress spurge</li> <li>- Field scabious</li> <li>- Giant hogweed</li> </ul>	<ul style="list-style-type: none"> <li>- Hoary cress</li> <li>- Japanese butterbur</li> <li>- Russian knapweed</li> <li>- Short-fringed knapweed</li> </ul>	<ul style="list-style-type: none"> <li>- Tree of heaven</li> <li>- Wild chervil</li> <li>- Wild parsnip</li> </ul>
<b>ANNUAL CONTROL</b> – Species with limited extent and/ or significant potential to spread. Management objective is to prevent further expansion into new areas with the ultimate goal of reducing the overall extent.		
<ul style="list-style-type: none"> <li>- Baby’s breath</li> <li>- Black knapweed (BC)</li> <li>- Bluweed</li> <li>- Bohemian knotweed</li> <li>- Brown knapweed</li> <li>- Himalayan blackberry</li> </ul>	<ul style="list-style-type: none"> <li>- Japanese knotweed</li> <li>- Leafy spurge (BC)</li> <li>- Poison hemlock</li> <li>- Policeman’s helmet</li> <li>- Rush skeletonweed (BC)</li> </ul>	<ul style="list-style-type: none"> <li>- Scotch broom</li> <li>- Scotch thistle</li> <li>- Teasel</li> <li>- Yellow archangel</li> <li>- Yellow flag iris</li> </ul>
<b>CONTAINMENT</b> – Species is established or with high potential for spread. Management objective is to prevent further expansion into new areas within the region through establishment of containment lines and identification of occurrences outside the line to control.		
<p><b>Contain to gardens:</b></p> <ul style="list-style-type: none"> <li>- Butterfly bush</li> <li>- Common periwinkle</li> <li>- English holly</li> <li>- English ivy</li> <li>- Garden yellow loosestrife</li> <li>- Goutweed</li> <li>- Mountain bluet</li> <li>- Myrtle spurge</li> </ul>	<p><b>Contain to gardens Cont’d:</b></p> <ul style="list-style-type: none"> <li>- Russian olive</li> <li>- Salt cedar/ Tamarisk</li> <li>- Siberian elm</li> </ul> <p><b>Contain to White Lake:</b></p> <ul style="list-style-type: none"> <li>- Fragrant water lily</li> </ul> <p><b>Contain to Crazy Creek RR:</b></p> <ul style="list-style-type: none"> <li>- Marsh plume thistle</li> </ul>	<p><b>Contain to west portion of IPMA (treat Seymour Arm and east portion of IPMA):</b></p> <ul style="list-style-type: none"> <li>- Common tansy</li> </ul> <p><b>Contain to Shuswap/ Mara/White Lake:</b></p> <ul style="list-style-type: none"> <li>- Eurasian water milfoil</li> </ul>
<b>MANAGEMENT</b> – Species is more widespread but may be of concern in specific situations with certain high values - e.g., conservation lands, specific agriculture crops. Management objective is to reduce the invasive species impacts locally or regionally, where resources are available.		
<ul style="list-style-type: none"> <li>- Bull thistle (BC)</li> <li>- Burdock</li> <li>- Canada thistle (BC)</li> <li>- Caraway</li> <li>- Cheatgrass/downy brome</li> <li>- Chicory</li> <li>- Common comfrey</li> <li>- Dalmatian toadflax (BC)</li> <li>- Diffuse knapweed (BC)</li> </ul>	<ul style="list-style-type: none"> <li>- Hoary alyssum</li> <li>- Hound’s tongue (BC)</li> <li>- Meadow buttercup</li> <li>- Meadow goat’s beard</li> <li>- Meadow knapweed (BC)</li> <li>- Nightshade</li> <li>- Orange hawkweed</li> <li>- Oxeye daisy</li> <li>- Purple loosestrife (BC)</li> </ul>	<ul style="list-style-type: none"> <li>- Queen Anne’s Lace</li> <li>- Spotted knapweed (BC)</li> <li>- St. John’s Wort (BC)</li> <li>- Sulphur cinquefoil</li> <li>- Western goat’s beard</li> <li>- Wormwood</li> <li>- Yellow hawkweed spp.</li> <li>- Yellow toadflax (BC)</li> </ul>
<b>INSUFFICIENT INFORMATION</b> – Species have insufficient information on their distribution, impacts, potential for spread and/or feasibility of control. Further information is required.		
<ul style="list-style-type: none"> <li>- Bachelor’s button</li> <li>- Black locust</li> <li>- Carpet burweed</li> <li>- Coltsfoot</li> <li>- Creeping buttercup</li> <li>- Curly leaf pondweed</li> <li>- Dame’s rocket</li> </ul>	<ul style="list-style-type: none"> <li>- Eyebright</li> <li>- Field bindweed</li> <li>- Flat peavine</li> <li>- Greater celandine</li> <li>- Green foxtail</li> <li>- Kochia</li> </ul>	<ul style="list-style-type: none"> <li>- Lady’s thumb</li> <li>- Russian thistle</li> <li>- Scentless chamomile (BC)</li> <li>- Sweet fennel</li> <li>- Wild four o’clock</li> <li>- Woolly vetch</li> </ul>

BC – biocontrol

Updated April 2026